

10/034, 985



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/20, 9/16, A23K 1/165 // (C12N 1/20, C12R 1:07)		A1	(11) International Publication Number: WO 97/33976
			(43) International Publication Date: 18 September 1997 (18.09.97)
(21) International Application Number: PCT/KR97/00040		(74) Agent: HUH, Sang, Hoon; Hyecheon Building, 13th floor, 831, Yeoksam-dong, Kangnam-ku, Seoul 135-792 (KR).	
(22) International Filing Date: 14 March 1997 (14.03.97)			
(30) Priority Data: 1996/6817 14 March 1996 (14.03.96) KR		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(71) Applicant (for all designated States except US): KOREA INSTITUTE OF SCIENCE AND TECHNOLOGY [KR/KR]; 39-1, Hawolgok-dong, Sungbuk-ku, Seoul 136-130 (KR).		Published With international search report. With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description. Date of receipt by the International Bureau: 25 June 1997 (25.06.97)	
(72) Inventors; and (75) Inventors/Applicants (for US only): OH, Tae Kwang [KR/KR]; Expo Apartment, 407-602, Junmin-dong, Yusung-ku, Daejeon 305-390 (KR). KIM, Hyung, Kwon [KR/KR]; Hanbit Apartment, 120-701, Uheun-dong, Yusung-ku, Daejeon 305-333 (KR). BAE, Kyung, Suk [KR/KR]; Clover Apartment, 115-1506, Dunsan-dong, Seo-ku, Daejeon 302-173 (KR). PARK, Young, Seo [KR/KR]; Hanul Apartment, 105-703, Sinsung-dong, Yusung-ku, Daejeon 305-345 (KR). KIM, Young, Ok [KR/KR]; 9/5 442-13, Jangjun-dong, Geumjung-ku, Pusan 609-390 (KR). CHE, Yang, Ung [KR/KR]; 546-10, Dogok-dong, Kangnam-ku, Seoul 135-270 (KR). LEE, Dong, Kuu [KR/KR]; Sindonga Apartment, 7-206, Subinggo-dong, Yongsan-ku, Seoul 140-240 (KR).			

(54) Title: DS11 (KCTC 0231BP), NOVEL BACILLUS SP. STRAIN AND NOVEL PHYTASE PRODUCED BY IT

(57) Abstract

This invention relates to a novel phytase produced from novel strain *Bacillus* sp. DS11 (KCTC 0231BP) and more precisely, to a novel strain *Bacillus* sp. DS11 and phytase enzyme enhancing the phosphate bioavailability present in grains supplied to monogastric animals.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**DS11 (KCTC 0231BP), NOVEL *BACILLUS SP.* STRAIN AND NOVEL
PHYTASE PRODUCED BY IT**

BACKGROUND OF THE INVENTION

5 **Field of the Invention**

This invention relates to a novel phytase produced from novel strain *Bacillus sp.* DS11 (KCTC 0231BP) and more precisely, to a novel strain *Bacillus sp.* DS11 and phytase enzyme enhancing the phosphate bioavailability present in grains supplied to monogastric animals.

10 **Description of the Prior Art**

Phytase is an enzyme to degrade phytic acid into phosphate and phosphate inositol. 50 to 70% of phosphate in grain used as livestock feeds exists in form of phytic acid but phytase is not present in monogastric animals such as hens and hogs, resulting in low phosphate availability.

15 Therefore undigested phytic acid(phytictaine) released to a water source became one of the serious environment contamination sources causing to eutrophication in small lakes or tides. With above consideration, because monogastric animals can't utilize phytic acid in their intestine phytic acid chelates to water due to chelation with a trace amount of minerals, amino
20 acids and vitamins which are very important to the metabolism of livestock. These formed water-insoluble, undigestable chelate complexes released to feces change the environmental ecosystem to induce a serious environmental pollution.

In view of these situations, the application of phytase into the livestock
25 feeds will reduce the supply of inorganic phosphate due to increase of phosphate bioavailability in livestock, leading to economic benefits, and improving the availability of phosphate, and other bioactive substances, leading to reduction of the environmental contamination.

By these reasons, the utilization of phytase in livestock is very important. A law regulating the amount of phosphate in animal waste was established in 1996 in Korea and in European countries it is already mandatory to add phytase in the feeds of animals.

5 The addition of phytase in the feeds may greatly improve the productivity of livestock by enhancing the availability of some bioactive substances(phosphate, calcium and zinc etc.) which, otherwise chelate with phytictaine and lose their activity.

As the result, the use of feeds containing phytase in livestock can
10 enhance the availability of feeds and reduce the environmental contamination caused by phosphate.

For the aforementioned benefits, intensive studies about phytase including the effects of phytase on animals (L.G. Young et al., 1993, X.G. Lei et al., 1994, Z. Mroz et al., 1994) have been performed mainly in Europe (A.H.J.
15 Ullah et al., 1994, K.C. Ehrich, 1994, C.S. Piddington, 1993). However, since phytase can cleave only a limited number of phosphates and it mostly produced by molds which have long growing period, it is uneconomical in terms of mass production. In addition, it is difficult to use the phytic acid as an additive for monogastric animals since it is unsuitable to their
20 physiological mechanism.

SUMMARY OF THE INVENTION

Therefore, the inventor et al. have identified a novel phytase-producing microorganism among hundred kinds of molds, Actinomycetes, bacteria, etc.,
25 obtained from soils and barns throughout the country, in an effort to produce phytase having excellent enzymatic potency and shorten the production period compared with the conventional phytase. Because of the high enzymatic potency of this novel microorganism and its physiological

suitability to use for monogastric animals and short period of kproduction compared to those of conventional enzymes the inventor et al. have judged that this enzyme has novelty and they have completed this invention.

The object of this invention is to provide novel strain *Bacillus sp.* DS11
5 (KCTC 0231BP) and phytase enzyme, which is suitable to use for monogastric animals with excellent properties and more shortened production period.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is electron microscope photograph (X 10,000) of *Bacillus sp.* DS11
10 (KCTC 0231BP) strain, a novel strain according to this invention,

Fig. 2 is a SDS-PAGE electrophoresis analysis of novel phytase produced by *Bacillus sp.* DS11 (KCTC 0231BP), A: precipitation of acetone, B: resource S, C: superose 12 HR 10/30

Fig. 3 is a graph measuring of novel phytase produced by *Bacillus sp.*
15 DS11 (KCTC 0231BP), a novel strain, in accordance with EXPERIMENTAL EXAMPLE 2 of this invention.

DETAILED DESCRIPTION OF THIS INVENTION

This invention relates to a novel phytase produced from novel strain
20 *Bacillus sp.* DS11 (KCTC 0231BP) and particularly, to a novel strain *Bacillus sp.* DS11 and phytase enzyme enhancing the phosphate bioavailability present in grains supplied to monogastric animals.

This invention is described in detail as set forth hereunder:

This invention relates to novel strain *Bacillus sp.* DS11(KCTC 0231BP).

25 Further, this invention includes its novel enzyme phytase whose N-terminal amino acid sequence is expressed by the following [Sequence table 1] under the following conditions:

Optimal temperature: 65 °C

Optimal pH: 7.0

Molecular weight: 43,000 dalton

Isoelectric point: 5.6

In addition, this invention includes a method to use said
5 microorganism as feed additives.

This invention is described in more detail as set forth hereunder:

This invention relates to a novel strain *Bacillus sp.* DS11 (KCTC 0231BP) and novel phytase produced from said strain. The procedure for isolating and identifying said novel microorganism is as follows:

10 **[Isolation of novel microorganisms]**

From several thousands of strains obtained from soils and in barns throughout the country, a strain was isolated which has excellent resolution in phytase screen plate containing 15g of D-glucose, 5 g of calcium phytate, 5g of NH_4NO_3 , 0.5 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g of KCl, 0.01 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g
15 $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ and agar 15 g at pH 7.0/ l.

[Identification of novel microorganism]

The morphological property of the strain isolated from the above procedure is as follows:

1) Morphological property

20 In the grams staining test of the strain cultivated in an optimal growth medium, the strain was proven to be gram-positive. Fig. 1, the picture of electron microscope shows that strain is a rod type of 0.8 ~ 1.8 μm in cell size.

In the growth test of the thermally inactivated cells at 80 °C, said strain has formed thermostable spores. Further, its catalase test in which it
25 shows the positive response demonstrates that it coincides with the morphological property of *Bacillus sp.*

The physiological property of said strain is as follows:

2) Physiological property

The test results of physiological property of the strain are represented in the following table 1. As shown in the table 1, it is assumed that strain is a facultative microorganism which may be grow in both aerobic and anaerobic states. With the difference of grown strains found at 50 °C and pH 5.7 compared with *Bacillus pantothenicus*, the strain of this invention is a mutant derived from *Bacillus pantothenicus*.

10

15

20

25

Table 1.

Property		<i>Bacillus</i> <i>pantotheniticus</i>	Strain of this invention
5	Catalase	+	+
	Anaerobic culture	+	+
	V-p. test	-	+
10	Acid-forming	glucose	+
		L-arabinose	+
		D-xylose	-
		D-mannitol	+
15	Hydrolysis	Casein	d
		Gelatin	+
		Starch	+
20	Synthesis	Indole	-
		Dihydroacetone	-
25	Growth pH	6.8	+
		5.7	+
	Concentration of growth salt (at NaCl)	2%	+
		5%	+
25	Growth temperature	7%	+
		10%	+
		5°C	-
		10°C	-
		30°C	+
25		40°C	+
		50°C	+

Note) +: positive. -: negative. d: different from species.

The chemical property of the strain obtained as above is as follows:

3) The chemical property of strain

After harvesting the strain, the test results of their various properties (e.g., G + C content, fatty acid composition, Murein Type and main melaquinone) are represented in the following table 2.

As shown in the above table 2, it is noted that the strain of this invention is similar to *Bacillus pantotheniticus* in terms of G + C content, murein type and main melaquinone including fatty acid of strain but it seems to difficult to make a judgement that both strains are the same species. Thus, it seems that the strain of this invention is a mutant derived from *Bacillus pantotheniticus*.

Table 2.

Property		<i>Bacillus pantotheniticus</i>	Strain of this invention
G + C content		36.9	45.6
Murein type		Mezo-DAP	Mezo-DAP
Main melaquinone		MK-7	MK-7
Fatty acid of strain (%)	14 : 0 ISO	4.71	1.96
	14 : 0	1.50	1.21
	15 : 0 ISO	19.34	18.72
	15 : 0 ANTEISO	37.95	38.51
	16 : 0 ISO	10.01	6.16
	16 : 0	9.77	9.77
	17 : 0 ISO	4.37	9.02
	17 : 0 ANTEISO	12.00	11.20
	18 : 0	x	2.03

By compiling above mentioned morphological, physiological and chemical properties, it is revealed that the strain of this invention belongs to *Bacillus sp.* from Bergys Manual of Systemic Bacteriology Vol. 2 (Williams & Wilkins Co., 1989). Therefore, the microorganism isolated was nominated
5 as *Bacillus sp.* DS11 and deposited on February 1, 1996, to Korea Research Institute of Bioscience and Biotechnology Korean Collection for type cultures, Korean Patent Strain Deposit Associations located in #52, Oun-dong, Yusong-ku, Taejeon 305-333, Republic of Korea, with an accession number KCTC 0231BP.

10 This invention is described in more detail by the following examples, but the claims are not limited to these examples.

EXAMPLE 1: Production of novel phytase

To produce phytase, a medium containing 6% wheat bran, NH_2NO_3 , 0.04%, 0.02% $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.0% casein hydrolysate, 0.05% KH_2PO_4 , 0.04%
15 K_2HPO_4 and 0.2% CaCl_2 , was adjusted to pH 6.5 and sterilized at 121 °C for 15 minutes.

Then, with the same medium composition, 1% seed culture cultivated in a flask at 37°C for 12 hours was inoculated into the medium in order to produce the enzyme.

20 The potency of produced enzyme was measured as follows: By using a substrate, comprising 2 mM phytic acid sodium salts and 2mM CaCl_2 in 0.1 M Tris buffer solution (pH 7.0), the enzyme was reacted at 37°C for 30 minutes to measure the amount of phosphoric acid generated. One unit of the enzymatic potency is equivalent to the enzymatic amount degrading 1 μ mol
25 of inorganic phosphate per 1 minute. As a result of said measurement, the novel enzyme has the enzymatic potency of 0.3 unit per protein mg. The maximum amount of enzyme in fermentation yielding 0.6 unit/mg was obtained when 30 l of strains, a working volume, was charged to a 50 l

fermenter and cultivated at 37 °C for 48 hours under the following conditions:
air influx - 0.8 vvm; stirring rate - 150 rpm. Compared with the enzymatic
production amount of 0.3 unit/mg when phytase enzyme derived from
Aspergillus ficuum is cultivated for 96 hours [Donna M. Gibson, 1987], it is well
5 observed that the novel organism of this invention has a production amount
twice higher.

**EXPERIMENTAL EXAMPLE 1: Measurement of molecular weight for novel
phytase**

2 1 of novel strain solution, cultivated according to the procedure as
10 described in said Example, was centrifuged for 15 minutes. Then, the
supernatant solution was saturated with 50% acetone to precipitate proteins
and the crude enzyme solution passed through dialysis membrane was
purified on a column containing phenyl sepharose CL-4B, Resource S and
superose 12 HR 10/30 (all of which are manufactured by Pharmacia, Sweden)
15 so as to isolate the phytase enzyme only. The analytical results by said
column are represented in the attached drawing Fig. 2. Phytase enzyme
was again on SDS-PAGE electrophoresis and the results show that the
molecular weight of phytase produced by a novel strain is 43,000 dalton and
isoelectric point is 5.6.

20 Further, the enzyme protein, so isolated, was used to determine the
sequence of N-terminal amino acid using Protein/peptide Sequencer (Applied
Biosystems, USA) and its results are represented in the following [Sequence
table].

The sequencing No. 1 of [Sequence table] is the sequence of N-terminal
25 amino acids of the phytase produced from a novel strain of this invention; the
sequencing No. 2 is the sequence of the N-terminal amino acids of the phytase
produced from *E. coli* [Arch. Biochem. Biophys. 303(1993)]; the sequencing No.
2 is the sequence of the N-terminal amino acids of the phytase produced from

Aspegillus ficuum [Prep. Biochem. 18(1988)].

From the results of [Sequence table], it is revealed that the phytase enzyme produced by *Bacillus* sp. DS11(KCTC 0231BP) of this invention is a novel enzyme.

5 **EXPERIMENTAL EXAMPLE 2: Activity and stability of novel phytase to temperature and pH**

The optimal temperature was proved to be 65℃ by using the enzyme isolated by the same method as described in the experimental example 2.

To measure the thermal stability, the enzyme was left at each temperature for
10 10 minutes so as to assess the residual activity. As shown in the attached drawing Fig. 3 (3-1), the results show that when Ca^{2+} was not added, the activity began to decrease at 40 ℃ but in case of adding 5mM calcium ion, the activity was stable up to 70℃ and 50% of the activity at 75℃ was sustained.

From the aforementioned results, a novel phytase of this invention may
15 be expected to a higher activity in the body of livestock. Therefore, it seems to be preferable that the feeds should be pelleted or extruded at more than 75℃ so as to use them as processing ones.

The results of phytase activity in the different conditions of pH are shown in the attached drawing Fig. 3 (3-2) and the optimal pH was 7.0.
20 Further, to measure the stability on pH, the enzyme was left at various conditions of pH for 1 hour so as to assess the residual activity of phytase. Even under the acidic condition of below pH 4, the enzyme was proved to have higher enzymatic activity and from these results, it is judged that the novel phytase of this invention may be extremely stable under the acidic
25 condition of stomach.

From the test results of the aforementioned temperatures and pH, it is noted that the novel phytase of this invention may be applicable as a feed additive of monogastric animals.

EXPERIMENTAL EXAMPLE 3: Influence of metal ion and inhibitor on enzyme activity

Influence of metal ion and inhibitor on enzyme activity is represented in the following table 3.

- 5 The results of table 3 show that the addition of 1mM EDTA inhibited the entire enzyme activity and when Cu^{2+} , Zn^{2+} and Mg^{2+} were added at the concentration of 5mM, about 50% of enzyme activity was reduced.

Table 3.

Additives	Concentration	
	1 mM	5 mM
Non-addition	100	100
CuCl_2	63	43
ZnCl_2	87	47
MgCl_2	95	49
MnCl_2	65	20
LiCl_2	95	100
HgCl_2	83	62
CaCl_2	99	116
RbCl_2	103	102
EDTA	7.5	7.6
PMSF	86	88

EXPERIMENTAL EXAMPLE 4: Influence of novel phytase added to feed for broiler on environmental contamination

To estimate the influence relation of availability and released amounts of phosphate when phytase is added, broilers were divided into three groups such as novel phytase group, soybean-extracted plant phytase group and mold

phytase group on the market (manufacturer: Sigma). Each of hatched-out 200 males of Avaachre broiler chicken was announced publicly for this experiment. The cultured medium containing phytase was ultrafiltrated and further concentrated at low temperature under vacuum and dried by lyophilizer. Each 500 unit of this lyophilized phytase was added to per kg of feed. The same amount of soybean phytase or mold phytase as above was also added to the feeds and the results were represented in the following table 4.

10 **Table 4**

Classification	Plant (soybean) phytase	Commercial mold phytase	Novel phytase
Phosphate contents in feeds (g/kg)	5.5	5.5	5.5
Intake of total feeds (g/kg)	2.7	2.7	2.7
15 Intake of phosphate (g/head)	15.1	15.2	15.1
Accumulation of phosphate (g/head)	7.1	7.8	8.8
Discharge of phosphate (g/head)	8.0	7.4	6.3
Absorption of phosphate (%)	47	51	58

20 The results of Table 4 revealed that when novel phytase was fed to monogastric animals, the phosphate contents released were lower than the soybean group or mold phytase group, since the phosphate availability in the former was higher than that of the latter; Namely, the activity of novel phytase was superior under acid-fast and acidic conditions and phytictaine
25 within the grains was degraded in the intestine of animals by the novel phytase in more effective manner.

[Sequence table]

Sequencing No.: 1

Sequencing length: 15

Sequencing form: amino acid

5 Shape: straight chain

Sequencing type: protein

Sequence: Ser-Asp-Pro-Tyr-His-Phe-Thr-Val-Asn-Ala-Ala-Xaa-Glu-Thr-Glu

Sequencing No.: 2

10 Sequencing length: 11

Sequencing form: amino acid

Shape: straight chain

Sequencing type: protein

Sequence: Ser-Glu-Pro-Glu-Leu-Lys-Leu-Glu-Ala-Val-Val

15

Sequencing No.: 3

Sequencing length: 12

Sequencing form: amino acid

Shape: straight chain

20 Sequencing type: protein

Sequence: Pro-Ala-Ser-Arg-x-Gin-Ser-Ser-Cys-Asp-Thr-Val

WHAT IS CLAIMED IS :

1. Strain *Bacillus sp.* DS11(KCTC 0231BP).
- 5 2. The method to use strain *Bacillus sp.* DS11(KCTC 0231BP) as feed additives.
3. The phytase produced by *Bacillus sp.* in claim 1 has the following characterization:
 - 10 Optimal temperature 65 °C
 - Optimal pH 7.0
 - Molecular weight 43,000 dalton
 - Isoelectric point 5.6
 - Sequence of N-terminal amino acid No. 1 based on [sequencing table]
- 15 4. The method to use a phytase produced by strain *Bacillus sp.* DS11(KCTC 0231BP) as feed additives according to claim 3.


FIG. 1



FIG. 2




FIG. 3-1



- ; optimal temperature
- ; stability in the non-existence of Ca^{2+}
- ; stability in the existence of 5mM Ca^{2+}

FIG.3-2



- ; optimal pH
- ; stability

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR 97/00040

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁶: C 12 N 1/20, 9/16; A 23 K 1/165 // (C 12 N 1/20; C 12 R 1:07)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁶: C 12 N 1/20, 9/16; A 23 K 1/165

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Patent Abstracts of Japan, Vol.18, No.257 (C-1200), 1994, Kokai-No. 6-38745 (ZENKOKU NOGYO KYODO KUMIAI RENGOKAI).	1-3
A	WO 94/19 471 A1 (THE FINNISH NATIONAL PUBLIC HEALTH INSTITUTE) 01 September 1994 (01.09.94), claims 1-15. -----	1-3

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 16 May 1997 (16.05.97)	Date of mailing of the international search report 02 June 1997 (02.06.97)
Name and mailing address of the ISA/ AT AUSTRIAN PATENT OFFICE Kohlmarkt 8-10 A-1014 Vienna Facsimile No. 1/53424/535	Authorized officer Wolf Telephone No. 1/53424/133

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/KR 97/00040

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WD A1 9419471	01-09-94	AU A1 61426/94	14-09-94
		CA AA 2156425	01-09-94
		CN A 1120852	17-04-96
		EP A1 686195	13-12-95
		FI A0 954010	25-08-95
		FI A 954010	25-10-95
		JP T2 8507207	06-08-96